

**Volunteer Biological Assessment Program
Stream Data Collection Report for the Saco Watershed
2022**



Green Mountain Conservation Group
P.O Box 95
236 Huntress Bridge Rd.
Effingham, NH 03882
(603) 539-1859
www.gmcg.org

Prepared by:

Grace Kennedy, GMCG Education and Outreach Assistant, Americorps Member
Tara Schroeder, GMCG Education Coordinator

ACKNOWLEDGEMENTS

The Green Mountain Conservation Group (GMCG) would like to acknowledge the continued training, guidance and support of Kayla Croteau of the NH Fish & Game Department and the many people and organizations who helped to make this program possible. Each year, volunteers and staff of GMCG collect the data presented in this annual report. GMCG would like to thank the teachers, aides, administrators, parents, care-givers, volunteers, students, bus drivers and local businesses who helped make the Volunteer Biological Assessment Program possible in 2022.

Volunteers & Staff

Jill Emerson, GMCG Water Quality Coordinator
Maddy Martin, GMCG Americorps Member
Tara Schroeder, GMCG Education Coordinator
Brittney Semone, GMCG Americorps Member
Moselle Spiller, GMCG Outreach Coordinator

Jeff Beavers
David Bellinghari
Blair Folts
Felicia Ledgard
Karen Vitek
Victor Vitek

Participating School Groups:

Effingham Elementary School
Freedom Elementary School
Ossipee Central School
Pine Tree Elementary School
Sandwich Central School

Community Partners:

The Other Store
Indian Mound Golf Course

This program is supported by:

The Alfred Quimby Fund
The Kendal C. and Anna Ham Charitable Foundation
New Hampshire Moose Plate Conservation Fund

Table Of Contents

Introduction	4
Methods	4
Sampling Sites	5
Data Collection	5
Macroinvertebrate Sorting and Identification	5
Biotic Index Computation	6
Supplementary Data	7
Water Quality Results	8
Future Recommendations	111
Appendix A: Site Maps and Pie Charts of Macroinvertebrates	12
Appendix B: Site Data Sheets	17
Appendix C: Microplastics Photos	212
Appendix C: Works Cited	27

Introduction

Since 1997, the New Hampshire Department of Environmental Services (DES) has conducted stream surveys to determine the health of aquatic ecosystems. As part of these efforts, DES developed a preliminary screening protocol for 1st through 4th order streams that is appropriate for volunteers to evaluate the biological condition of aquatic macroinvertebrate communities. The goals of the protocol are as follows:

- To educate the public about water quality issues as interpreted through biological assessments;
- To build a constituency of individuals who will practice sound water quality management at the local level; and
- To build public support for water quality protection.

Since 2006, Green Mountain Conservation Group (GMCG) has collaborated with DES, NH Fish & Game Department and local volunteers and schools for the Volunteer Biological Assessment Program (VBAP). While NH DES no longer oversees the program “due to lack of staff support” according to David Neils, Chief Aquatic Pollution Biologist, GMCG continues to offer the VBAP program to schools in the Ossipee Watershed for educational purposes following the recommendation of NH DES and NH Fish & Game Department. In recent years, the program has expanded to additional schools and sites outside of the Ossipee Watershed, including parts of the Saco River Watershed in Conway, NH and Porter, ME.

In 2022, GMCG worked with students and teachers from five local schools across the Saco Watershed in New Hampshire and Maine to collect, sort, and analyze macroinvertebrates at five sites. This report contains the results of that sampling with additional data analysis to compare sites across the watershed and over time since some of these sites are year-round sampling sites for GMCG’s RIVERS program (Regional Interstate Volunteers for the Ecosystems and Rivers of Saco). As monitoring continues, data will continue to be evaluated and analyzed for any trends or water quality issues.

Methods

Prior to sampling, a training was held in September at GMCG’s Conservation Center with NH Fish & Game for new teachers and volunteers in the program. A training session was also held at each school for students during a scheduled classroom period and consisted of the following components: macroinvertebrate identification skills; biomonitoring and macroinvertebrate tolerance levels; macroinvertebrate sampling protocols; and an introduction to watersheds; riparian ecosystems; and aquatic food chains. Additionally, student volunteers were trained to collect and record supplementary data for physical and chemical parameters of the rivers/streams, including: habitat assessments; stream measurements; pH; conductivity; temperature; and dissolved oxygen.

In 2022, GMCG staff and AmeriCorps members also incorporated microplastics sampling as part of the program for the second year. Students were introduced in the classroom to the concept of microplastics in the environment and were able to collect water samples during the field day which were analyzed in GMCG’s water quality resource center. Students collected a sample by placing a plankton net in an area with moving current. The net was left in the current for at least five minutes, and was then retrieved, a water sample was collected and transferred to

a glass storage container. In the GMCG lab, 100 mL of each sample was filtered through a Buchner funnel with a 0.45 μm filter. After filtering, the filter paper was transferred to a glass plate for observation. The first sample was examined by students in the classroom using a white light microscope. 10 mL of 10 μg Nile Red solution was added to the second sample. This sample was then examined in the GMCG water quality lab using a microscope equipped with blue light and an orange filter. Under these lighting conditions, nonpolar particles (i.e. plastic) treated with Nile Red will appear fluorescent.

Sampling Sites

All five sampling sites were accessible, wadeable, approximately 200 feet in length, 1st through 4th order streams and contained appropriate sampling habitat (at least one riffle, one pool, and one run with mixed cobble substrate).

Additional sites are sampled from spring through fall for GMCG's Regional Interstate Volunteers for the Ecosystems and Rivers of Saco (RIVERS) program, and have been periodically sampled by school groups in the past for VBAP. While some schools sample their adopted VBAP site annually, some are on an every-other-year schedule due to combined classes. Sampling was scheduled throughout September and October and required three to four hours per site.

Data Collection

The data collection at each site and along the stream followed the same protocol. The protocol was as follows:

- A 500-micron mesh kicknet was placed perpendicular to stream flow and held firmly against the streambed with the opening of the net facing upstream to promote macro invertebrate collection.
- A collector would disturb the sample area (1/5 m^2) upstream of the net for a total of 60 seconds (30 second hand-scrub followed by a 30 second kick).
- The kicknet was carefully lifted out of the water and the contents of the net were emptied into a shallow container with a small amount of water. All organisms remaining on the net were carefully removed and added to the sample.
- The same process was repeated four additional times with each sample collected further upstream (spanning 200 feet). Collectively, active sampling time approximated five minutes within one square meter area at each sampling station.

Macroinvertebrate Sorting and Identification

For approximately 60 minutes, student volunteers, teachers and/or staff removed macroinvertebrates from the selected portion of the sample with spoons or pipettes and placed them into separate containers according to common attributes. After sorting, specimens were identified to various coarse taxonomic groups, and the number of macroinvertebrates within each taxonomic group was identified, calculated, and recorded, see Table 1. Students were assisted by GMCG staff and/or trained volunteers with the process of identifying the macroinvertebrates in the sample.

Table 1. Total Macroinvertebrates Found Across the Watershed in 2022. For individual site macroinvertebrate counts see Appendix A.

Order	Common Name	Number of Macroinvertebrates
Ephemeroptera	Mayfly nymph	389
Plecoptera	Stonefly nymph	80
Trichoptera	Caddisfly larvae	362
Odonata	Dragonfly larvae	8
	Damselfly nymph	1
Diptera	Black fly larvae	8
	Midge larvae	9
	True flies	13
Megaloptera	Alderfly	0
	Hellgrammite	13
Coleoptera	Riffle Beetle	22
	Water Penny	2
	Beetle/Beetle like	0
Other	Crayfish	0
	Snails	0
	Aquatic Worms	15
	Scuds	0
	Sowbug	0
	Leech	0
	Water Mites	18
	Planaria	12

Biotic Index Computation

Biotic scores are based on pollution tolerance values ranging from 0 to 9 and are assigned to individual taxonomic groups. More tolerant groups have higher tolerance values and less tolerant groups have lower values. A standardized computational worksheet was used to compute the biotic scores for each sample site (stream/river). Taxonomic-specific biotic scores for individual samples were computed by multiplying the number of individual organisms by their respective tolerance value; summing the taxonomic-specific biotic scores; and then dividing the sum by the total number of individuals identified in the respective sample. Final biotic scores correspond to three interim narrative categories:

- Excellent (0 to 3.5)
- Good (3.5 to 4.8)
- Fairly Poor (greater than 4.8).

Table 2. Biotic Scores of Sampling Sites. Table 2 highlights the biotic score and the narrative category for each sampling site tested in the Ossipee Watershed. Four of the five scores fell in the “Excellent” category while one fell in the “Good” category.

School Group	Date	Location	Town	Total Number of Macros	Biotic Score	Water Quality Score
Effingham Elementary	9/20/22	South River	Parsonsfield, ME	332	3.68	Good
Sandwich Central School	9/24/22	Cold River	Sandwich, NH	293	3.09	Excellent
Freedom Elementary	9/26/22	Lovell River	Ossipee, NH	80	3.08	Excellent
Pine Tree Elementary	9/23/22	Swift River	Conway, NH	121	3.32	Excellent
Ossipee Central School	9/28/22	Swift River	Tamworth, NH	126	3.29	Excellent




Figure 1. Water Quality Scores. The graph shows changes in water quality scores over time at the sites sampled in 2022 and in years past.

Supplementary Data

The water chemistry and physical parameters of the stream were recorded. Physical parameters recorded included width/depth of the stream, canopy cover, observations of nearby erosion or human influence, pH, conductivity and temperature. A multi-parameter submersible

Each water quality probe was used to collect pH, dissolved oxygen, conductivity and temperature data (see Table 3). Microplastics samples were collected using a standard plankton net.

Water Quality Results

Basic water quality data were collected at each of the sampling locations. The data included chemical parameters, physical parameters, and calculating a biotic score for each of the sampling sites. Macroinvertebrate samples from each site were evaluated using the VBAP biotic score index utilizing taxa-specific tolerance values. A cumulative biotic score for all sites and individual site-specific biotic scores were computed. The average biotic score for all sites was 3.44 and corresponds to the “Excellent” narrative category. See Table 2 for biotic scores for each of the individual sampling sites. Overall, mayfly nymphs were the most dominant taxon 389 (41%), followed by caddisfly larvae 362 (38%) and stonefly nymphs 80 (8.4%). Together, these three taxa comprised nearly 87.4% of all the individuals collected and are some of the least tolerant taxonomic groups. In completing the sampling effort, volunteers collected and staff identified 952 macroinvertebrates (See Table 1).

Table 3. Physical & Chemical Parameters of Sampling Sites

Physical & Chemical Parameters	pH	Dissolved Oxygen (mg/L)	Conductivity (µS/cm)	Temperature (°C)
Normal Range/Optimal Value	6.0-8.0	5 and above	Below 100	Below 21°C for trout/mayfly nymphs
South River	6.065	8.075	63.7	13.4
Cold River	6.00	11.48	26.7	8.2
Lovell River	6.00	10.29	21.28	13
Swift River (Conway)	5.5	10.6	40.1	12.5
Swift River (Tamworth)	6.00	10.8	34.4	11

Dissolved oxygen, and temperature were conducive to supporting aquatic life at all sites. pH at all sites and particularly the Swift River in Conway, were slightly low or acidic. pH values in streams are affected naturally by the bedrock or streambed material and the flow of groundwater into the stream. Streams with an underlying bedrock of limestone for example have a greater ability to buffer acidity in a stream and raise pH. According to the Saco River Corridor Commission’s (SRCC) [2021 report](#):


pH influences chemical and biological processes that occur in water and are essential for aquatic organisms. The ability of aquatic organisms to complete a life cycle greatly diminishes as pH falls below 5.0 or exceeds 9.0. Levels below 5.5 can severely limit growth and reproduction in fish, as is the case with brook trout in New England streams. Low pH can also allow toxic elements and compounds such as heavy metals to become mobile and available for uptake by aquatic plants and animals, which in turn can cause deformities in fish and produce conditions that are toxic to aquatic life. These low pH

levels can be due to naturally occurring conditions, such as the influence of tannic and humic acids from decaying plants in wetlands. Low pH can also be influenced by industrial pollution in the form of atmospheric deposition of nitric and sulfuric acids in acid rain. The discharge of wastewater from treatment plants can also affect natural pH.”

The SRCC 2021 report shows that the Swift River site in Conway has a median pH level that falls within Maine’s state standards (above 6.5, based on 110 samples). In addition, as stated in GMCG’s RIVERS program data analysis, the waterbodies in the Ossipee Watershed are mostly underlain by granite, which does not have the ability to buffer acidity, and “surface waters in this area tend to be naturally around a pH of 6.5”.


Evidence of microplastics was found at each of the five sampling sites from samples collected by the students using plankton nets (see Figure 2).

Figure 2. Evidence of microplastics was found at each of the five sites. From left to right these images show a microbead, a microfragment, and a microfilament of plastic collected from Swift River in Tamworth.



Conductivity is a measure of the concentration of dissolved salts in water. Conductivity levels are not a concern to the EPA (Environmental Protection Agency) unless they rise above 500 $\mu\text{S}/\text{cm}^2$ for rural areas or 1500 $\mu\text{S}/\text{cm}^2$ for urban areas, according to Jill Emerson, GMCG’s Water Quality Coordinator. For reference, seawater has a Specific Conductance of 55,000 μS . NH DES states that conductivity levels in freshwater bodies across the state are rising, in general, mostly due to road salting, faulty septic systems and urban/agricultural runoff (NHDES, 2004). While conductivity levels across all VBAP sampling sites remain low, conductivity levels at South River are slightly elevated and data shows a steady increase in the median annual conductivity in the South River over the past twenty years. Simple trend line analysis of historical data taken through GMCG’s RIVERS program demonstrates that conductivity levels at this site are slightly increasing over time, likely due to road salting activities (see Figure 3).

Figure 3. South River chemical and physical data trends from 2002 - 2022. The analysis shows a steady increase in the median annual conductivity in the South River over the past twenty years.



To bring this issue of slowly rising conductivity levels to the attention of Ossipee Watershed towns, GMCG has co-hosted workshops such as Green SnowPro with UNH T2 Center, NH DES and NH Department of Transportation to help train area road agents and plow drivers on efficient use/spreading of salt to help educate those on the front lines about the importance of not over-salting our freshwater ecosystems. New Hampshire's freshwater lakes, streams and groundwater are becoming saltier each year. The leading cause is the 400,000 tons of road salt applied every winter to our Interstates, state highways, town roads and other surfaces. Salt is toxic to aquatic life and plants, it corrodes bridges and vehicles, and when it accumulates in drinking water it jeopardizes human health.

Green Mountain Conservation Group launched the Salt Responsibly campaign in January 2022 to inform New Hampshire residents about the



harm caused by road salts and to provide guidance on ways to reduce the amount of salt that is contaminating New Hampshire's waterways and water supply.

The Salt Responsibly campaign is not about casting blame. We recognize that safety is the first priority of those responsible for winter road maintenance, and until the day that practical alternatives are widely available, road salt use will continue. What we seek is a broader public understanding of the problem and to inspire everyone to make changes that will reduce the amount of salt that is damaging our environment, threatening our health, and hurting our economy.


For more detailed water quality analyses on the water bodies in the Ossipee Watershed, please see the 10 Year Water Quality Report (GMCG, 2015) or town reports as recent as 2021: <http://www.gmcg.org/research/water-quality-program-data/>. The SRCC's water quality report of data from 2001 through 2020, published in 2021, is also available: https://srcc-maine.org/wp-content/uploads/2022/01/SRCC_2020WQManalysis_FBE_FinalDraft.pdf.

Future Recommendations

The documentation of the invertebrate communities by volunteers using the VBAP protocol during the fall of 2022 in the Ossipee Watershed marked the seventeenth year of 'screening' efforts to evaluate the status of aquatic communities. Sampling efforts included five sites in the Saco Watershed. Macroinvertebrates are widely used as indicators of water quality that can show the effects of multiple pollutants over time. It is important to recognize that the results obtained from the VBAP protocol are not intended to represent formal water quality assessments, but rather, a basic indicator of aquatic community condition. Ultimately, the results from the VBAP for 2022 build upon the efforts conducted by GMCG, NH DES and NH Fish & Game to establish reference sites in the state, compare sites across the state, and provide watershed education to youth. The results of the program serve as a basis for further monitoring and management practices to be put into use throughout the watershed. Schools also use their VBAP data for the Trout in the Classroom program to make sure the water quality conditions and macroinvertebrates at their release sites are adequate to support Eastern brook trout. These data are in fact required by NH Fish and Game in order to issue schools permits for the program.


The Ossipee Watershed has a reputation for having great water quality overall and the VBAP results validate this statement. It is recommended that GMCG continue to work with schools in the Ossipee Watershed and continue to monitor the water quality of the local streams and rivers. Biotic scores for sites sampled in 2022 ranged from good to excellent, and continued monitoring will be essential to track any water quality changes over time. The long-term health of surface waters will depend on preventing potential sources of contamination from entering water bodies and using best management practices that reduce or prevent adverse impacts from human activities, such as road salting. Sites where human activities appear to be impacting water quality should be monitored closely for potential sources of contamination, and road salting alternatives should be sought. In addition, continued monitoring for microplastics pollution is recommended as a means of educating students about the importance of reducing plastic use and the local impacts of this global problem.

Appendix A: Site Maps and Pie Charts of Macroinvertebrates



(Above) Total macroinvertebrates found at South River in Parsonsfield, ME(Below)







(Above) Total Macroinvertebrates found at Cold River in Sandwich, NH (Below)




2022 Total Macroinvertebrates Freedom Elementary (Lovell River)





(Above) Total macroinvertebrates found at Lovell River in Ossipee, NH (Below)






(Above) Total macroinvertebrates found at Swift River in Tamworth, NH (Below)





(Above) Total macroinvertebrates found at Swift River in Conway, NH (Below)



Appendix B: Site Data Sheets

South River, Parsonsfield, ME

Order	Common Name	Value	*	Found	=	Score	Score	Category
Ephemeroptera	Mayfly Nymph	3	*	56	=	168	3.2	Excellent
Plecoptera	Stonefly Nymph	1	*	9	=	9		
Trichoptera	Caddisfly Larvae	4	*	41	=	164		
Odonata	Dragonfly Nymph	3	*	4	=	12		
	Damselfly Nymph	7	*	1	=	7		
Diptera	Black fly larvae	7	*	0	=	0		
	Midge larvae	6	*	2	=	12		
	Most True flies	4	*	1	=	4		
Megaloptera	Alderfly	4	*	0	=	0		
	Hellgrammite	0	*	8	=	0		
Coleoptera	Riffle beetle	4	*	18	=	72		
	Water Penny	4	*	0	=	0		
	Beetle & Beetle-like	7	*	0	=	0		
Others	Crayfish	6	*	0	=	0		
	Snails	7	*	0	=	0		
	Aquatic Worms	8	*	0	=	0		
	Scuds	8	*	0	=	0		
	Sowbugs	7	*	0	=	0		
	Clams and Mussels	7	*	0	=	0		
Totals				140		448	3.2	Excellent

Cold River, Sandwich, NH

Order	Common Name	Value	*	Found	=	Score	Score	Category
Ephemeroptera	Mayfly Nymph	3	*	161	=	483		
Plecoptera	Stonefly Nymph	1	*	14	=	14		
Trichoptera	Caddisfly Larvae	4	*	91	=	364		
Odonata	Dragonfly Nymph	3	*	1	=	3		
	Damselfly Nymph	7	*	0	=	0		
Diptera	Black fly larvae	7	*	5	=	35		
	Midge larvae	6	*	0	=	0		
	Most True flies	4	*	8	=	32		
Megaloptera	Alderfly	4	*	0	=	0		
	Hellgrammite	0	*	1	=	0		
Coleoptera	Riffle beetle	4	*	4	=	16		
	Water Penny	4	*	2	=	8		
	Beetle & Beetle-like	7	*	0	=	0		
Others	Crayfish	6	*	0	=	0		
	Snails	7	*	0	=	0		
	Aquatic Worms	8	*	0	=	0		
	Scuds	8	*	0	=	0		
	Sowbugs	7	*	0	=	0		
	Clams and Mussels	7	*	0	=	0		
Totals				287		955	3.32	Excellent

Lovell River, Ossipee, NH

Order	Common Name	Value	*	Found	=	Score	Score	Category
Ephemeroptera	Mayfly Nymph	3	*	45	=	135		
Plecoptera	Stonefly Nymph	1	*	15	=	15		
Trichoptera	Caddisfly Larvae	4	*	8	=	32		
Odonata	Dragonfly Nymph	3	*	0	=	0		
	Damselfly Nymph	7	*	0	=	0		
Diptera	Black fly larvae	7	*	0	=	0		
	Midge larvae	6	*	4	=	24		
	Most True flies	4	*	2	=	8		
Megaloptera	Alderfly	4	*	0	=	0		
	Hellgrammite	0	*	2	=	0		
Coleoptera	Riffle beetle	4	*	0	=	0		
	Water Penny	4	*	0	=	0		
	Beetle & Beetle-like	7	*	0	=	0		
Others	Crayfish	6	*	0	=	0		
	Snails	7	*	0	=	0		
	Aquatic Worms	8	*	4	=	32		
	Scuds	8	*	0	=	0		
	Sowbugs	7	*	0	=	0		
	Clams and Mussels	7	*	0	=	0		
	Water Mites	4	*	0	=	0		
Totals				80		246	3.25	Excellent

Swift River, Conway, NH




Order	Common Name	Value	*	Found	=	Score	Score	Category
Ephemeroptera	Mayfly Nymph	3	*	56	=	168		
Plecoptera	Stonefly Nymph	1	*	9	=	9		
Trichoptera	Caddisfly Larvae	4	*	41	=	164		
Odonata	Dragonfly Nymph	3	*	2	=	6		
	Damselfly Nymph	7	*	0	=	0		
Diptera	Black fly larvae	7	*	3	=	21		
	Midge larvae	6	*	1	=	6		
	Most True flies	4	*	2	=	8		
Megaloptera	Alderfly	4	*	0	=	0		
	Hellgrammite	0	*	2	=	0		
Coleoptera	Riffle beetle	4	*	0	=	0		
	Water Penny	4	*	5	=	20		
	Beetle & Beetle-like	7	*	0	=	0		
Others	Crayfish	6	*	0	=	0		
	Snails	7	*	0	=	0		
	Aquatic Worms	8	*	0	=	0		
	Scuds	8	*	0	=	0		
	Sowbugs	7	*	0	=	0		
	Clams and Mussels	7	*	0	=	0		
Totals				121		402	3.32	Excellent

Swift River, Tamworth, NH



Order	Common Name	Value	*	Found	=	Score	Score	Category
Ephemeroptera	Mayfly Nymph	3	*	47	=	141	3.29	Excellent
Plecoptera	Stonefly Nymph	1	*	30	=	30		
Trichoptera	Caddisfly Larvae	4	*	30	=	120		
Odonata	Dragonfly Nymph	3	*	1	=	3		
	Damselfly Nymph	7	*	0	=	0		
Diptera	Black fly larvae	7	*	0	=	0		
	Midge larvae	6	*	2	=	12		
	Most True flies	4	*	0	=	0		
Megaloptera	Alderfly	4	*	0	=	0		
	Hellgrammite	0	*	0	=	0		
Coleoptera	Riffle beetle	4	*	0	=	0		
	Water Penny	4	*	0	=	0		
	Beetle & Beetle-like	7	*	0	=	0		
Others	Crayfish	6	*	0	=	0		
	Snails	7	*	0	=	0		
	Aquatic Worms	8	*	11	=	88		
	Scuds	8	*	0	=	0		
	Sowbugs	7	*	0	=	0		
	Clams and Mussels	7	*	0	=	0		
	Water Mites	4	*	5	=	20		
Totals				126		414	3.29	Excellent

Appendix C: Microplastics Photos



South River, Parsonsfield, ME






Cold River, Sandwich, NH






Lovell River, Ossipee, NH



Swift River, Conway, NH



Swift River, Tamworth, NH



Appendix C: Works Cited

FB Environmental & Saco River Corridor Commission (2021, December). *Saco River Corridor Commission 2020 Water Quality Analysis*. https://srcc-maine.org/wp-content/uploads/2022/01/SRCC_2020WQManalysis_FBE_FinalDraft.pdf

Green Mountain Conservation Group (2015, March). *Ossipee Watershed 10-Year Water Quality Report*. <http://www.gmcg.org/research/10-year-water-quality-report/>.

New Hampshire Department of Environmental Services (2004). *Special Topics Article: Conductivity is on the rise in New Hampshire's lakes and ponds: What is causing the increase and what can be done?* https://www.des.nh.gov/organization/commissioner/pip/publications/wd/documents/2004_special_topic_conductivity.pdf